

Northwest Invasive Plant Council Priority Plant List

The Northwest Invasive Plant Council (NWIPC) keeps a dynamic Invasive Plant Priority List for the region. This list is used by NWIPC staff to develop work plans based on the region, climate and partners needs. This list is broken up into Invasive Plant Management Areas (IPMA's) due to the diversity of plants and climate across the region.

Definitions

<p><u>Provincial Early Detection Rapid Response (Provincial EDRR)</u></p>	<p>Invasive Plants within the Provincial EDRR Process are not present in B.C., or are present but limited in extent and being actively managed with the goal of eradication from B.C.. These plants pose a significant threat to B.C.'s environment, economy, and/or human health.</p> <p>All Provincial EDRR species on the NWIPC list are identified in red with an (*)</p> <p>Provincial Priority Invasive Species List</p>
<p><u>Regional Early Detection Rapid Response (REDRR)</u></p>	<p>Early Detection Rapid Response species pose a significant threat and is new to the region. These species exist in the region/IPMA, but with very limited distribution. Eradication is feasible. REDRR Species will generally be less than 10 plants in abundance in any given IPMA.</p> <p><u>Management Approach:</u> Eradication is the goal. These species are the highest priority for planned control programs.</p>
<p><u>High Priority</u></p>	<p>These species are abundant in certain portions of the IPMA/region but have not yet infested all potential habitats. Plants that have passed the threshold of REDRR (generally greater than 10 plants per IPMA) will be moved to High Priority.</p> <p><u>Management Approach:</u> High risk/impact; limited population with significant potential to spread in the region. Contain to prevent further expansion. If the population is low enough, the objective will be to control every site annually.</p>
<p><u>Medium/Low Priority</u></p>	<p>These are widespread species that are beyond landscape-level control and/or have relatively low impact. Species are moved from High Priority to Medium/Low Priority when the committee decides that the current capacity of the NWIPC can no longer feasibly manage all of the REDRR, and high priority species with the given budget. Some species will be selected to have the priority decreased to low based on risk and eradication potential.</p> <p><u>Management Approach:</u> Medium risk/impact; limited distribution – broader population distribution with potential to spread further in a region. The goal of management efforts for these species is to protect site-specific values or assets (e.g., corridors of spread, agricultural land, etc.). Land managers may choose to treat these species at sites they deem valuable to protect based on specific land management objectives.</p>
<p><u>Containment Areas</u></p>	<p>NWIPC Containment Areas are established when high risk species become so abundant that there are not enough resources to feasibly control the population in an area. The goal of containments lines is that treatments can be focused outside the area and work inwards to control the spread of the species.</p>

NWIPC Regional Invasive Plant List

Invasive Species historically known to be in the NWIPC operating area

Common name	Genus species
Annual hawksbeard	<i>Crepis tectorum</i>
Baby's-breath	<i>Gypsophila paniculata</i>
Bachelor's button	<i>Centura cyanus</i>
Bighead knapweed	<i>Centaurea macrocephala</i>
Bishop's Goutweed	<i>Aegopodium podagraria</i>
*Black henbane	<i>Hyoscyamus niger</i>
Black knapweed	<i>Centaurea nigra</i>
Bladder campion	<i>Silene cucubalus</i>
Blueweed	<i>Echium vulgare</i>
Bohemian knotweed	<i>Fallopia x bohemica</i>
Brown knapweed	<i>Centaurea jacea</i>
Bull thistle	<i>Cirsium vulgare</i>
Bur chervil	<i>Anthriscus caucalis</i>
Butterfly bush	<i>Buddleia davidii</i>
Camel thorn	<i>Vachellia erioloba</i>
Canada thistle	<i>Cirsium arvense</i>
Caraway	<i>Carum carvi</i>
Chicory	<i>Cichorium intybus</i>
Chilean tarweed	<i>Madia sativa</i>
Coltsfoot	<i>Tussilago farfara</i>
Common bugloss	<i>Anchusa officinalis</i>
Common burdock	<i>Arctium minus</i>
Common comfrey	<i>Symphytum officinale L.</i>
Common periwinkle	<i>Littorina littorea</i>
Common tansy	<i>Tanacetum vulgare</i>
Creeping buttercup	<i>Ranunculus repens</i>
Cudweed	<i>Gamochoeta spp</i>
Curled dock	<i>Rumex crispus L</i>
Cutleaf blackberry	<i>Rubus laciniatus</i>
Cypress spurge	<i>Euphorbia cyparissias</i>
Dalmatian toadflax	<i>Linaria dalmatica</i>
Dames rocket	<i>Hesperis matronalis</i>
Diffuse knapweed	<i>Centaurea diffusa</i>
Dodder	<i>Cuscuta cephalanthi</i>
English Holly	<i>Ilex aquifolium</i>
English ivy	<i>Hedera helix</i>
European bittersweet/ climbing nightshade	<i>Solanum dulcamara</i>
Eyebright	<i>Euphrasia officinalis</i>
Fernleaf yarrow	<i>Achillea filipendulina</i>
Field Bindweed	<i>Convolvulus arvensis</i>

Field scabious	<i>Knautia arvensis</i>
Garden yellow loosestrife	<i>Lysimachia vulgaris</i>
Giant Hogweed	<i>Heracleum mantegazzianum</i>
Giant knotweed	<i>Fallopia sachalinensis</i>
Gorse	<i>Ulex europaeus</i>
Greater celandin	<i>Chelidonium majus</i>
Greater knapweed	<i>Centaurea scabiosa</i>
Green foxtail	<i>Setaria viridis</i>
Groundsel	<i>Senecio vulgaris</i>
Hairy cat's-ear	<i>Hypochaeris radicata</i>
Heart-podded hoarycress	<i>Lepidium / Cardaria draba</i>
Himalayan blackberry	<i>Rubus discolor</i>
Himalayan knotweed	<i>Persicaria wallichii</i>
Hoary alyssum	<i>Berteroa incana</i>
Hounds-tongue	<i>Cynoglossum officinale</i>
*Japanese butterbur	<i>Petasites japonicus</i>
Japanese knotweed	<i>Fallopia japonica</i>
Kochia	<i>Bassia / Kochia scoparia</i>
Large yellow loosestrife	<i>Lysimachia punctata L</i>
Leafy spurge	<i>Euphorbia esula</i>
Marsh plume thistle	<i>Cirsium palustre</i>
Meadow buttercup	<i>Ranunculus acris</i>
Meadow goat's-beard	<i>Tragopogon pratensis</i>
Meadow knapweed	<i>Centaurea debeauxii</i>
Milk thistle	<i>Silybum marianum</i>
Mossy stone crop	<i>Sedum acre</i>
Mountain bluet	<i>Centaurea montana</i>
*Mouse ear hawkweed	<i>Pilosella officinarum/ Hieracium pilosella</i>
Mullein	<i>Verbascum thapsus</i>
Night flowering catchfly	<i>Silene noctiflora</i>
Nodding thistle	<i>Carduus nutans</i>
Old man's beard	<i>Clematis vitalba</i>
Orange hawkweed	<i>Hieracium aurantiacum</i>
Oxeye daisy	<i>Chrysanthemum leucanthemum</i>
Plumeless thistle	<i>Carduus acanthoides</i>
Poison Hemlock	<i>Conium maculatum</i>
Policeman's helmet / Himalayan balsam	<i>Impatiens glandulifera</i>
Purple loosestrife	<i>Lythrum salicaria</i>
Queen Anne's lace / wild carrot	<i>Daucus carota</i>
Redroot amaranth / rough pigweed	<i>Amaranthus retroflexus</i>
Russian knapweed	<i>Acroptilon repens</i>
Russian thistle	<i>Salsola kali</i>
Scentless chamomile	<i>Matricaria maritima</i>
Scotch Broom	<i>Cytisus scoparius</i>
Scotch thistle	<i>Onopordum acanthium</i>

Sheep sorrel	<i>Rumex acetosella</i>
Shepherd's purse	<i>Capsella bursa-pastoris</i>
Sow thistle species	<i>Sonchus spp.</i>
Spanish bluebells	<i>Hyacinthoides hispanica</i>
Spotted Hawkweed	<i>Hieracium maculatum</i>
Spotted knapweed	<i>Centaurea stoebe</i>
St. John's wort/Saint John's wort/ Goatweed	<i>Hypericum perforatum</i>
Sulphur cinquefoil	<i>Potentilla recta</i>
Sweet fennel	<i>Foeniculum vulgare</i>
Tansy ragwort	<i>Senecio jacobaea</i>
Tartary buckwheat	<i>Fagopyrum tataricum</i>
Western goat's beard	<i>Tragopogon dubius</i>
Whiplash hawkweed	<i>Hieracium flagellare</i>
White cockle	<i>Silene latifolia</i>
Wild buckwheat	<i>Polygonum convolvulus</i>
Wild carrot/Queen annes lace	<i>Daucus carota</i>
Wild Chervil	<i>Anthriscus sylvestris</i>
Wild mustard	<i>Sinapis arvensis</i>
Wormwood/absinthium	<i>Artemisia absinthium</i>
Yellow archangel	<i>Lamium galeobdolon</i>
Yellow flag iris	<i>Iris pseudacorus</i>
*Yellow floating heart	<i>Nymphoides peltata</i>
Yellow hawkweeds	<i>Hieracium spp.</i>
Yellow loosestrife	<i>Lysimachia vulgaris</i>
Yellow toadflax	<i>Linaria vulgaris</i>

Bulkley IPMA

CONTAINMENT LINE PRESENT

*Provincial EDRR

Regional EDRR	High Priority	Lower Priority
Black knapweed (<i>Centaurea nigra</i>)	Common tansy (<i>Tanacetum vulgare</i>)	Annual hawksbeard (<i>Crepis tectorum</i>)
Blueweed (<i>Echium vulgare</i>)	Chicory (<i>Cichorium intybus</i>)	Bachelor's button (<i>Centaurea cyanus</i>)
Brown knapweed (<i>Centaurea jacea</i>)	Field scabious (<i>Knautia arvensis</i>)	Bladder campion (<i>Silene vulgaris</i>)
Chilean tarweed (<i>Madia sativa</i>)	Himalayan balsam (<i>Impatiens glandulifera</i>)	Bull thistle (<i>Cirsium vulgare</i>)
Common bugloss (<i>Anchusa officinalis</i>)	Mountain bluet (<i>Centaurea montana</i>)	Burdock species (<i>Arctium spp.</i>)
Cypress spurge (<i>Euphorbia cyparissias</i>)	St. John's wort/ Goatweed (<i>Hypericum perforatum</i>)	Canada thistle (<i>Cirsium arvense</i>)
Diffuse knapweed (<i>Centaurea diffusa</i>)	Yellow flag iris (<i>Iris pseudacorus</i>)	Common comfrey (<i>Symphytum officinale</i>)
Goutweed/bishops weed (<i>Aegopodium podagraria</i>)		Creeping buttercup (<i>Ranunculus repens</i>)
Greater knapweed (<i>Centaurea scabiosa</i>)		Curled dock (<i>Rumex crispus</i>)
Hoary alyssum (<i>Berteroa incana</i>)		Dalmation toadflax (<i>Linaria dalmatica</i>)
Japanese knotweed (<i>Reynoutria japonica</i>)		Dame's rocket (<i>Hesperis matronalis</i>)
Leafy spurge (<i>Euphorbia esula</i>)		Groundsel (<i>Senecio vulgaris</i>)
Marsh plume thistle (<i>Cirsium palustre</i>)		Meadow buttercup (<i>Ranunculus acris</i>)
Nodding thistle		Meadow salsify / goats-beard (<i>Tragopogon pratensis</i>)
Spotted hawkweed (<i>Hieracium maculatum</i>)		Mullein (<i>Verbascum thapsus</i>)
Sulphur cinquefoil (<i>Potentilla recta</i>)		Old man's beard (<i>Clematis vitalba</i>)
Tansy ragwort (<i>Senecio jacobaea</i>)		Orange hawkweed (<i>Hieracium aurantiacum</i>)
*Yellow floating heart (<i>Nymphoides peltata</i>)		Oxeye daisy (<i>Leucanthemum vulgare</i>)
		Scentless chamomile (<i>Tripleurospermum inodorum</i>)
		Sow thistle species (<i>Sonchus spp.</i>)
		Western salsify/ goat's-beard (<i>Tragopogon dubius</i>)
		Wormwood (<i>Artemisia absinthium</i>)
		Yellow/common toadflax (<i>Linaria vulgaris</i>)
		Yellow hawkweed species (<i>Hieracium / Pilosella spp.</i>)

Lakes District IPMA

CONTAINMENT LINE PRESENT

*Provincial EDRR

Regional EDRR	High Priority	Lower Priority
Baby's breath (<i>Gypsophila paniculata</i>)	Common tansy (<i>Tanacetum vulgare</i>)	Annual hawksbeard (<i>Crepis tectorum</i>)
Bighead knapweed (<i>Centaurea macrocephala</i>)	Mountain bluet (<i>Centaurea montana</i>)	Bull thistle (<i>Cirsium vulgare</i>)
Black knapweed (<i>Centaurea nigra</i>)	St. John's wort/ Goatweed (<i>Hypericum perforatum</i>)	Burdock species (<i>Arctium spp.</i>)
Cypress spurge (<i>Euphorbia cyparissias</i>)	Spotted knapweed (<i>Centaurea stoebe</i>)	Canada thistle (<i>Cirsium arvense</i>)
Diffuse knapweed (<i>Centaurea diffusa</i>)		Chicory (<i>Cichorium intybus</i>)
Field scabious (<i>Knautia arvensis</i>)		Common comfrey (<i>Symphytum officinale</i>)
Kochia (<i>Bassia / Kochia scoparia</i>)		Curled dock (<i>Rumex crispus</i>)
Leafy Spruge (<i>Euphorbia esula</i>)		Dalmation toadflax (<i>Linaria dalmatica</i>)
Meadow knapweed (<i>Centaurea debeauxii</i>)		Field bindweed (<i>Convolvulus arvensis</i>)
Queen Anne's lace/Wild carrot (<i>Daucus carota</i>)		Meadow buttercup (<i>Ranunculus acris</i>)
		Meadow salsify / goats-beard (<i>Tragopogon pratensis</i>)
		Mullein (<i>Verbascum thapsus</i>)
		Orange hawkweed (<i>Hieracium aurantiacum</i>)
		Oxeye daisy (<i>Leucanthemum vulgare</i>)
		Scentless chamomile (<i>Tripleurospermum inodorum</i>)
		Sheep sorrel (<i>Rumex acetosella</i>)
		Sow thistle species (<i>Sonchus spp.</i>)
		Western salsify/ goat's-beard (<i>Tragopogon dubius</i>)
		White cockle (<i>Silene latifolia / Lychnis alba</i>)
		Yellow hawkweed species (<i>Hieracium / Pilosella spp.</i>)
		Yellow/common toadflax (<i>Linaria vulgaris</i>)

Nechako IPMA

CONTAINMENT LINE PRESENT

*Provincial EDRR

Regional EDRR	High Priority	Lower Priority
Baby's breath (<i>Gypsophila paniculata</i>)	Brown knapweed (<i>Centaurea jacea</i>)	Annual hawksbeard (<i>Crepis tectorum</i>)
Blueweed (<i>Echium vulgare</i>)	Common tansy (<i>Tanacetum vulgare</i>)	Bachelor's button (<i>Centaurea cyanus</i>)
Camel thorn (<i>Alhagi maurorum</i>)	Field scabious (<i>Knautia arvensis</i>)	Bladder campion (<i>Silene vulgaris</i>)
Chilean tarweed (<i>Madia sativa</i>)	Himalayan balsam (<i>Impatiens glandulifera</i>)	Bull thistle (<i>Cirsium vulgare</i>)
Japanese knotweed (<i>Reynoutria japonica</i>)	Leafy spurge (<i>Euphorbia esula</i>)	Burdock species (<i>Arctium spp.</i>)
Meadow knapweed (<i>Centaurea debeauxii</i>)	Marsh plume thistle (<i>Cirsium palustre</i>)	Canada thistle (<i>Cirsium arvense</i>)
Nodding thistle (<i>Carduus nutans</i>)	Mountain bluet (<i>Centaurea montana</i>)	Chicory (<i>Cichorium intybus</i>)
Sulphur cinquefoil (<i>Potentilla recta</i>)	Spotted knapweed (<i>Centaurea stoebe</i>)	Curled dock (<i>Rumex crispus</i>)
Tansy ragwort (<i>Senecio jacobaea</i>)	St. John's wort/ Goatweed (<i>Hypericum perforatum</i>)	Dalmation toadflax (<i>Linaria dalmatica</i>)
Whiplash hawkweed (<i>Hieracium flagellare</i>)		Dame's rocket (<i>Hesperis matronalis</i>)
		Field bindweed (<i>Convolvulus arvensis</i>)
		Meadow buttercup (<i>Ranunculus acris</i>)
		Meadow salsify / goats-beard (<i>Tragopogon pratensis</i>)
		Mullein (<i>Verbascum thapsus</i>)
		Orange hawkweed (<i>Hieracium aurantiacum</i>)
		Oxeye daisy (<i>Leucanthemum vulgare</i>)
		Scentless chamomile (<i>Tripleurospermum inodorum</i>)
		Sow thistle species (<i>Sonchus spp.</i>)
		Western salsify/ goat's-beard (<i>Tragopogon dubius</i>)
		White cockle (<i>Silene latifolia</i> / <i>Lychnis alba</i>)
		Wormwood (<i>Artemisia absinthium</i>)
		Yellow/common toadflax (<i>Linaria vulgaris</i>)
		Yellow hawkweed species (<i>Hieracium</i> / <i>Pilosella spp.</i>)

Prince George IPMA

CONTAINMENT LINE PRESENT

*Provincial EDRR

Regional EDRR	High Priority	Lower Priority
Baby's breath (<i>Gypsophila paniculata</i>)	Brown knapweed (<i>Centaurea jacea</i>)	Annual hawkbeard (<i>Crepis tectorum</i>)
Bighead knapweed (<i>Centaurea macrocephala</i>)	Common tansy (<i>Tanacetum vulgare</i>)	Bachelor's button (<i>Centaurea cyanus</i>)
*Black henbane (<i>Hyoscyamus niger</i>)	Cypress spurge (<i>Euphorbia cyparissias</i>)	Bladder campion (<i>Silene vulgaris</i>)
Blueweed (<i>Echium vulgare</i>)	Diffuse knapweed (<i>Centaurea diffusa</i>)	Bull thistle (<i>Cirsium vulgare</i>)
Bohemian knotweed (<i>Reynoutria x bohemica</i>)	Field scabious (<i>Knautia arvensis</i>)	Burdock species (<i>Arctium spp.</i>)
Chilean tarweed (<i>Madia sativa</i>)	Himalayan balsam (<i>Impatiens glandulifera</i>)	Canada thistle (<i>Cirsium arvense</i>)
Common bugloss (<i>Anchusa officinalis</i>)	Hoary alyssum (<i>Berteroa incana</i>)	Chicory (<i>Cichorium intybus</i>)
Garden yellow loosestrife (<i>Lysimachia vulgaris</i>)	Japanese knotweed (<i>Reynoutria japonica</i>)	Common comfrey (<i>Symphytum officinale</i>)
Greater knapweed (<i>Centaurea scabiosa</i>)	Leafy spurge (<i>Euphorbia esula</i>)	Creeping buttercup (<i>Ranunculus repens</i>)
Green foxtail (<i>Setaria viridis</i>)	Marsh plume thistle (<i>Cirsium palustre</i>)	Curled dock (<i>Rumex crispus</i>)
Kochia (<i>Bassia / Kochia scoparia</i>)	Mountain bluet (<i>Centaurea montana</i>)	Dalmation toadflax (<i>Linaria dalmatica</i>)
Yellow/ spotted loosestrife (<i>Lysimachia punctata</i>)	Nodding thistle (<i>Carduus nutans</i>)	Dame's rocket (<i>Hesperis matronalis</i>)
Meadow knapweed (<i>Centaurea debeauxii</i>)	Spotted knapweed (<i>Centaurea stoebe</i>)	Eyebright (<i>Euphrasia nemorosa</i>)
Purple loosestrife (<i>Lythrum salicaria</i>)	St. John's wort/ Goatweed (<i>Hypericum perforatum</i>)	Field bindweed (<i>Convolvulus arvensis</i>)
Queen Anne's lace/Wild carrot (<i>Daucus carota</i>)	Sulphur cinquefoil (<i>Potentilla recta</i>)	Groundsel (<i>Senecio vulgaris</i>)
Russian knapweed (<i>Rhaponticum / Acroptilon repens</i>)	Whiplash hawkweed (<i>Hieracium flagellare</i>)	Hairy cat's-ear (<i>Hypochaeris radicata</i>)
Russian thistle (<i>Salsola tragus / kali</i>)		Meadow buttercup (<i>Ranunculus acris</i>)
Scotch broom (<i>Cytisus scoparius</i>)		Meadow salsify / goats-beard (<i>Tragopogon pratensis</i>)
Scotch thistle (<i>Onopordum acanthium</i>)		Mullein (<i>Verbascum thapsus</i>)
Tansy ragwort (<i>Senecio jacobaea</i>)		Night flowering catchfly (<i>Silene noctiflora</i>)
Yellow flag iris (<i>Iris pseudacorus</i>)		Orange hawkweed (<i>Hieracium aurantiacum</i>)
		Oxeye daisy (<i>Leucanthemum vulgare</i>)
		Plumeless thistle (<i>Carduus acanthoides</i>)
		Redroot amaranth/ rough pigweed (<i>Amaranthus retroflexus</i>)
		Scentless chamomile (<i>Tripleurospermum inodorum</i>)
		Shepherd's purse (<i>Capsella bursa-pastoris</i>)
		Sow thistle species (<i>Sonchus spp.</i>)
		Western salsify/ goat's-beard (<i>Tragopogon dubius</i>)
		Wormwood (<i>Artemisia absinthium</i>)
		Yellow hawkweed species (<i>Hieracium / Pilosella spp.</i>)
		Yellow/common toadflax (<i>Linaria vulgaris</i>)

Robson Valley IPMA

CONTAINMENT LINE PRESENT

*Provincial EDRR

Regional EDRR	High Priority	Lower Priority
Bighead knapweed (<i>Centaurea macrocephala</i>)	Baby's-breath (<i>Gypsophila paniculata</i>)	Annual hawksbeard (<i>Crepis tectorum</i>)
Blueweed (<i>Echium vulgare</i>)	Common tansy (<i>Tanacetum vulgare</i>)	Bladder campion (<i>Silene vulgaris</i>)
Brown knapweed (<i>Centaurea jacea</i>)	Cypress spurge (<i>Euphorbia cyparissias</i>)	Bull thistle (<i>Cirsium vulgare</i>)
Field scabious (<i>Knautia arvensis</i>)	Diffuse knapweed (<i>Centaurea diffusa</i>)	Canada thistle (<i>Cirsium arvense</i>)
Heart-podded hoary cress (<i>Lepidium / Cardaria draba</i>)	Hoary alyssum (<i>Berteroa incana</i>)	Chicory (<i>Cichorium intybus</i>)
Meadow knapweed (<i>Centaurea debeauxii</i>)	Marsh plume thistle (<i>Cirsium palustre</i>)	Common comfrey (<i>Symphytum officinale</i>)
Nodding thistle (<i>Carduus nutans</i>)	Mountain bluet (<i>Centaurea montana</i>)	Creeping buttercup (<i>Ranunculus repens</i>)
Himalayan balsam (<i>Impatiens glandulifera</i>)	Spotted knapweed (<i>Centaurea stoebe</i>)	Curled dock (<i>Rumex crispus</i>)
Sulphur cinquefoil (<i>Potentilla recta</i>)	St. John's wort/ Goatweed (<i>Hypericum perforatum</i>)	Dalmation toadflax (<i>Linaria dalmatica</i>)
Tansy ragwort (<i>Senecio jacobaea</i>)	Whiplash hawkweed (<i>Hieracium flagellare</i>)	Dame's rocket (<i>Hesperis matronalis</i>)
Yellow archangel (<i>Lamiaeastrum galeobdolon</i>)		Meadow buttercup (<i>Ranunculus acris</i>)
		Meadow salsify / goats-beard (<i>Tragopogon pratensis</i>)
		Mullein (<i>Verbascum thapsus</i>)
		Night flowering catchfly (<i>Silene noctiflora</i>)
		Orange hawkweed (<i>Hieracium aurantiacum</i>)
		Oxeye daisy (<i>Leucanthemum vulgare</i>)
		Scentless chamomile (<i>Tripleurospermum inodorum</i>)
		Sow thistle species (<i>Sonchus spp.</i>)
		Western salsify/ goat's-beard (<i>Tragopogon dubius</i>)
		White cockle (<i>Silene latifolia / Lychnis alba</i>)
		Wild mustard (<i>Sinapis arvensis</i>)
		Wormwood (<i>Artemisia absinthium</i>)
		Yellow hawkweed species (<i>Hieracium / Pilosella spp.</i>)
		Yellow/common toadflax (<i>Linaria vulgaris</i>)

Skeena IPMA

CONTAINMENT LINE PRESENT

*Provincial EDRR

Regional EDRR	High Priority	Lower Priority
Baby's-breath (<i>Gypsophila paniculata</i>)	Brown knapweed (<i>Centaurea jacea</i>)	Annual hawksbeard (<i>Crepis tectorum</i>)
Black knapweed (<i>Centaurea nigra</i>)	Common tansy (<i>Tanacetum vulgare</i>)	Bull thistle (<i>Cirsium vulgare</i>)
Blueweed (<i>Echium vulgare</i>)	Diffuse knapweed (<i>Centaurea diffusa</i>)	Burdock species (<i>Arctium spp.</i>)
Bohemian knotweed (<i>Reynoutria x bohemica</i>)	English ivy (<i>Hedera helix</i>)	Canada thistle (<i>Cirsium arvense</i>)
Chicory (<i>Cichorium intybus</i>)	Garden yellow loosestrife (<i>Lysimachia vulgaris</i>)	Common comfrey (<i>Symphytum officinale</i>)
Chilean tarweed (<i>Madia sativa</i>)	Giant knotweed (<i>Reynoutria / Fallopia sachalinensis</i>)	Common periwinkle (<i>Vinca minor</i>)
Common bugloss (<i>Anchusa officinalis</i>)	Himalayan knotweed (<i>Persicaria wallichii</i>)	Creeping buttercup (<i>Ranunculus repens</i>)
Coltsfoot (<i>Tussilago farfara</i>)	Japanese knotweed (<i>Reynoutria japonica</i>)	Curled dock (<i>Rumex crispus</i>)
English holly (<i>Ilex aquifolium</i>)	Yellow/ spotted loosestrife (<i>Lysimachia punctata</i>)	Dalmation toadflax (<i>Linaria dalmatica</i>)
Field scabious (<i>Knautia arvensis</i>)	Marsh plume thistle (<i>Cirsium palustre</i>)	Dame's rocket (<i>Hesperis matronalis</i>)
Greater knapweed (<i>Centaurea scabiosa</i>)	Mountain bluet (<i>Centaurea montana</i>)	Goutweed / Bishop's weed (<i>Aegopodium podagraria</i>)
Hounds-tongue (<i>Cynoglossum officinale</i>)	Himalayan balsam (<i>Impatiens glandulifera</i>)	Groundsel (<i>Senecio vulgaris</i>)
*Japanese butterbur (<i>Petasites japonicus</i>)	Scotch broom (<i>Cytisus scoparius</i>)	Meadow buttercup (<i>Ranunculus acris</i>)
*Mouse ear hawkweed (<i>Pilosella officinarum</i> Vaill.)	Spotted knapweed (<i>Centaurea stoebe</i>)	Mullein (<i>Verbascum thapsus</i>)
Nodding thistle (<i>Carduus nutans</i>)	St. John's wort/ Goatweed (<i>Hypericum perforatum</i>)	Night flowering catchfly (<i>Silene noctiflora</i>)
Purple loosestrife (<i>Lythrum salicaria</i>)	Tansy ragwort (<i>Senecio jacobaea</i>)	Orange hawkweed (<i>Hieracium aurantiacum</i>)
Sulphur cinquefoil (<i>Potentilla recta</i>)	Yellow archangel (<i>Lamiastrum galeobdolon</i>)	Oxeye daisy (<i>Leucanthemum vulgare</i>)
Whiplash hawkweed (<i>Hieracium flagellare</i>)	Yellow flag iris (<i>Iris pseudacorus</i>)	Scentless chamomile (<i>Tripleurospermum inodorum</i>)
Wild Chervil (<i>Anthriscus sylvestris</i>)		Sheep sorrel (<i>Rumex acetosella</i>)
		Sow thistle species (<i>Sonchus spp.</i>)
		Spanish bluebells (<i>Hyacinthoides hispanica</i>)
		Tartary buckwheat (<i>Fagopyrum tataricum</i>)
		Western salsify/ goat's-beard (<i>Tragopogon dubius</i>)
		Wormwood (<i>Artemisia absinthium</i>)
		Yellow hawkweed species (<i>Hieracium / Pilosella spp.</i>)
		Yellow/common toadflax (<i>Linaria vulgaris</i>)

Stikine IPMA

CONTAINMENT LINE PRESENT

*Provincial EDRR

Regional EDRR	High Priority	Lower Priority
Black knapweed (<i>Centaurea nigra</i>)	Common tansy (<i>Tanacetum vulgare</i>)	Annual hawksbeard (<i>Crepis tectorum</i>)
Burdock species (<i>Arctium spp.</i>)	Mountain bluet (<i>Centaurea montana</i>)	Bladder campion (<i>Silene vulgaris</i>)
Dalmation toadflax (<i>Linaria dalmatica</i>)	Orange hawkweed (<i>Hieracium aurantiacum</i>)	Bull thistle (<i>Cirsium vulgare</i>)
Diffuse knapweed (<i>Centaurea diffusa</i>)	Scentless chamomile (<i>Tripleurospermum inodorum</i>)	Canada thistle (<i>Cirsium arvense</i>)
Field scabious (<i>Knautia arvensis</i>)	Spotted knapweed (<i>Centaurea stoebe</i>)	Common comfrey (<i>Symphytum officinale</i>)
Marsh plume thistle (<i>Cirsium palustre</i>)	St. John's wort/ Goatweed (<i>Hypericum perforatum</i>)	Creeping buttercup (<i>Ranunculus repens</i>)
Nodding thistle (<i>Carduus nutans</i>)		Curled dock (<i>Rumex crispus</i>)
Whiplash hawkweed (<i>Hieracium flagellare</i>)		Dodder (<i>Cuscuta spp.</i>)
		Groundsel (<i>Senecio vulgaris</i>)
		Meadow buttercup (<i>Ranunculus acris</i>)
		Oxeye daisy (<i>Leucanthemum vulgare</i>)
		Sheep sorrel (<i>Rumex acetosella</i>)
		Sow thistle species (<i>Sonchus spp.</i>)
		Western salsify/ goat's-beard (<i>Tragopogon dubius</i>)
		Yellow hawkweed species (<i>Hieracium / Pilosella spp.</i>)
		Yellow/common toadflax (<i>Linaria vulgaris</i>)

Haida Gwaii IPMA

CONTAINMENT LINE PRESENT

*Provincial EDRR

Regional EDRR	High Priority	Lower Priority
Bighead knapweed (<i>Centaurea macrocephala</i>)	Bohemian knotweed (<i>Reynoutria x bohemica</i>)	Annual sow thistle (<i>Sonchus oleraceus</i>)
Butterfly bush (<i>Buddleja davidii</i>)	Common tansy (<i>Tanacetum vulgare</i>)	Bull thistle (<i>Cirsium vulgare</i>)
Cutleaf blackberry (<i>Rubus laciniatus</i>)	Gorse (<i>Ulex europaeus</i>)	Burdock species (<i>Arctium spp.</i>)
Cypress spurge (<i>Euphorbia cyparissias</i>)	Himalayan blackberry (<i>Rubus armeniacus</i>)	Canada thistle (<i>Cirsium arvense</i>)
Diffuse knapweed (<i>Centaurea diffusa</i>)	Himalayan knotweed (<i>Persicaria wallichii</i>)	Common comfrey (<i>Symphytum officinale</i>)
English holly (<i>Ilex aquifolium</i>)	Japanese knotweed (<i>Reynoutria japonica</i>)	Curled dock (<i>Rumex crispus</i>)
English ivy (<i>Hedera helix</i>)	Scotch broom (<i>Cytisus scoparius</i>)	Eyebright (<i>Euphrasia nemorosa</i>)
Garden yellow loosestrife (<i>Lysimachia vulgaris</i>)	Tansy ragwort (<i>Senecio jacobaea</i>)	Groundsel (<i>Senecio vulgaris</i>)
Himalayan balsam (<i>Impatiens glandulifera</i>)	Yellow archangel (<i>Lamium galeobdolon</i>)	Hairy cat's-ear (<i>Hypochaeris radicata</i>)
Marsh plume thistle (<i>Cirsium palustre</i>)	Yellow flag iris (<i>Iris pseudacorus</i>)	Nightshade (<i>Solanum spp.</i>)
Mountain bluet (<i>Centaurea montana</i>)		Oxeye daisy (<i>Leucanthemum vulgare</i>)
Orange hawkweed (<i>Hieracium aurantiacum</i>)		Perennial sow thistle (<i>Sonchus arvensis</i>)
Spotted hawkweed (<i>Hieracium maculatum</i>)		Wild buckwheat (<i>Fallopia convolvulus</i>)
Spotted knapweed (<i>Centaurea stoebe</i>)		Wormwood (<i>Artemisia absinthium</i>)
St. John's wort/ Goatweed (<i>Hypericum perforatum</i>)		Yellow/common toadflax (<i>Linaria vulgaris</i>)
Yellow hawkweed species (<i>Hieracium / Pilosella spp.</i>)		